

# Rosetta Scoring (or Energy) Function

**Benjamin K. Mueller**

Postdoc, Meiler Lab


Rosetta Workshop

Nov 2018

# Rosetta Scoring (or Energy) Function

... or how models are evaluated in  
Rosetta

# Evaluating Models in Rosetta



An algorithm must assign a quantitative number to a model to decide whether it is “good” or “bad”

# Important Note

$$\Delta E_{\text{total}} = \sum_i w_i E_i(\theta_i, a_i)$$

Energy is currently given in:  
“kcal/mol”

Previously in: “REU”


High Energy = Bad

Low Energy = Good

# Evaluating Models in Rosetta




-400 kcals/mol



-350 kcals/mol

# Score is Central to Monte Carlo Selection




# Metropolis Criterion

If  $E_{\text{new}} < E_{\text{old}}$  : Accept new structure

If  $E_{\text{new}} > E_{\text{old}}$  :

- Pick a random number  $p(0, 1)$

- if  $e^{-(E_{\text{new}} - E_{\text{old}})/kBT} > p$ , accept new structure




# Rosetta Combines Physics-Based and Knowledge-Based Potentials to Build the Energy Function

# Rosetta Combines Physics-Based and Knowledge-Based Potentials to Build the Energy Function

Lennard-Jones Potential

$$\sum_{i < j} \left[ \frac{A_{ij}}{r_{ij}^{12}} - \frac{B_{ij}}{r_{ij}^6} \right]$$

van der Waals Energy




# Rosetta Combines Physics-Based and Knowledge-Based Potentials to Build the Energy Function

Lennard-Jones Potential


$$\sum_{i < j} \left[ \frac{A_{ij}}{r_{ij}^{12}} - \frac{B_{ij}}{r_{ij}^6} \right]$$

van der Waals Energy



Ramachandran Plot

Statistical mining of  
Protein Databank (PDB)



# The Score Function is a Weighted Linear Combination of Individual Score Terms



Lennard-Jones Potential

$$\sum_{i < j} \left[ \frac{A_{ij}}{r_{ij}^{12}} - \frac{B_{ij}}{r_{ij}^6} \right]$$


Ramachandran Plot

Statistical mining of  
Protein Databank (PDB)

$$\text{Energy} = w_1 * \text{term}_1 + w_2 * \text{term}_2 + w_3 * \text{term}_3 + \dots$$



# Rosetta is a Residue-Centric Scoring Function



# Rosetta is a Residue-Centric Scoring Function


One Body	Two Body	Whole Body
Backbone - p_aa_pp - rama_prepro	Lennard-Jones - fa_atr - fa_rep	Radius of Gyration - rg
Side Chain - fa_dun - yhh_planarity	Solvation - fa_sol	Contact Order - co
Reference - ref	Hydrogen Bond - hbond_lr_bb - hbond_sr_bb - hbond_bb_sc - hbond_sc	Structure Alignment - hs_pair - ss_pair - sheet

\* NOTE: Not all score terms are listed here

# Rosetta has 2 score function modes

## Low resolution (or centroid) mode:


Reduced atom representation (centroid)  
Simple energy function  
Aggressively search conformational space



`database/chemical/residue_type_sets/centroid`

## High resolution (or full atom) mode:

Full atom (FA)  
More sophisticated energy function  
“Local” search of conformational (and sequence) space



`database/chemical/residue_type_sets/fa_standard`

# Breakdown of Full Atom Score Terms – REF2015

term	description	weight	units	ref(s)
fa_atr	attractive energy between two atoms on different residues separated by a distance $d$	1.0	kcal/mol	5, 6
fa_rep	repulsive energy between two atoms on different residues separated by a distance $d$	0.55	kcal/mol	5, 6
fa_intra_rep	repulsive energy between two atoms on the same residue separated by a distance $d$	0.005	kcal/mol	5, 6
fa_sol	Gaussian exclusion implicit solvation energy between protein atoms in different residues	1.0	kcal/mol	36
lk_ball_wtd	orientation-dependent solvation of polar atoms assuming ideal water geometry	1.0	kcal/mol	50, 71
fa_intra_sol	Gaussian exclusion implicit solvation energy between protein atoms in the same residue	1.0	kcal/mol	36
fa_elec	energy of interaction between two nonbonded charged atoms separated by a distance $d$	1.0	kcal/mol	50
hbond_lr_bb	energy of short-range hydrogen bonds	1.0	kcal/mol	38, 49
hbond_sr_bb	energy of long-range hydrogen bonds	1.0	kcal/mol	38, 49
hbond_bb_sc	energy of backbone–side-chain hydrogen bonds	1.0	kcal/mol	38, 49
hbond_sc	energy of side-chain–side-chain hydrogen bonds	1.0	kcal/mol	38, 49
dslf_fa13	energy of disulfide bridges	1.25	kcal/mol	49
rama_prepro	probability of backbone $\phi$ , $\psi$ angles given the amino acid type	(0.45 kcal/mol)/ $kT$	$kT$	50, 51
p_aa_pp	probability of amino acid identity given backbone $\phi$ , $\psi$ angles	(0.4 kcal/mol)/ $kT$	$kT$	51
fa_dun	probability that a chosen rotamer is native-like given backbone $\phi$ , $\psi$ angles	(0.7 kcal/mol)/ $kT$	$kT$	52
omega	backbone-dependent penalty for cis $\omega$ dihedrals that deviate from $0^\circ$ and trans $\omega$ dihedrals that deviate from $180^\circ$	(0.6 kcal/mol)/AU	AU <sup>a</sup>	72
pro_close	penalty for an open proline ring and proline $\omega$ bonding energy	(1.25 kcal/mol)/AU	AU	51
yhh_planarity	sinusoidal penalty for nonplanar tyrosine $\chi_3$ dihedral angle	(0.625 kcal/mol)/AU	AU	49
ref	reference energies for amino acid types	(1.0 kcal/mol)/AU	AU	1, 51

<sup>a</sup>AU = arbitrary units.

What you'll actually see in your  
output...

# Output Score Table

Found in output score table (score.sc) and at the end of every output pdb (S\_0001.pdb)

```
# All scores below are weighted scores, not raw scores.
#BEGIN_POSE_ENERGIES_TABLE dock_01_girk_1212_jon_ML297_0500.pdb
label fa_atr fa_rep fa_sol fa_intra_rep fa_elec total
weights 0.8 0.4 0.6 0.004 0.42 NA
pose -5400.73 638.748 2428.12 13.5145 -302.036 -4267.34
GLN:NtermProteinFull_1 -1.02843 0.12113 0.83426 0.00595 -0.05882 0.54065
ARG_2 -5.46203 0.44148 4.02307 0.01303 -1.13134 -3.04715
PHE_3 -5.25196 0.29047 1.51945 0.02354 -0.38757 -1.99927
VAL_4 -4.7212 0.38219 2.30519 0.01101 -0.33608 -4.69107
ASP_5 -6.18239 0.68119 4.3013 0.02667 -1.21805 -4.77894
LYS_6 -3.70129 0.51352 2.90682 0.00832 -0.90685 -3.25917
ASN_7 -2.97985 0.42136 1.9979 0.00362 -0.17245 -2.08876
GLY_8 -2.75353 0.28458 1.44949 1e-05 -0.20107 -4.00172
ARG:CtermProteinFull_1294 -2.34298 0.24806 1.73197 0.01216 -0.06632 -0.34395
```

# Output Score Table

## Score Terms:

```
# All scores below are weighted scores, not raw scores.
#BEGIN POSE ENERGIES TABLE dock 01_girk 1212_jon ML297_0500.pdb
label fa atr fa rep fa sol fa intra rep fa elec total
weights 0.8 0.4 0.6 0.004 0.42 NA
pose -5400.73 638.748 2428.12 13.5145 -302.036 -4267.34
GLN:NtermProteinFull_1 -1.02843 0.12113 0.83426 0.00595 -0.05882 0.54065
ARG_2 -5.46203 0.44148 4.02307 0.01303 -1.13134 -3.04715
PHE_3 -5.25196 0.29047 1.51945 0.02354 -0.38757 -1.99927
VAL_4 -4.7212 0.38219 2.30519 0.01101 -0.33608 -4.69107
ASP_5 -6.18239 0.68119 4.3013 0.02667 -1.21805 -4.77894
LYS_6 -3.70129 0.51352 2.90682 0.00832 -0.90685 -3.25917
ASN_7 -2.97985 0.42136 1.9979 0.00362 -0.17245 -2.08876
GLY_8 -2.75353 0.28458 1.44949 1e-05 -0.20107 -4.00172
ARG:CtermProteinFull_1294 -2.34298 0.24806 1.73197 0.01216 -0.06632 -0.34395
```

# Output Score Table

## Weights:

```
# All scores below are weighted scores, not raw scores.  
#BEGIN POSE ENERGIES TABLE dock 01_girk 1212_jon ML297_0500.pdb  
label fa_atr fa_rep fa_sol fa_intra_rep fa_elec total  
weights 0.8 0.4 0.6 0.004 0.42 NA  
pose -5400.73 638.748 2428.12 13.5145 -302.036 -4267.34  
GLN:NtermProteinFull_1 -1.02843 0.12113 0.83426 0.00595 -0.05882 0.54065  
ARG_2 -5.46203 0.44148 4.02307 0.01303 -1.13134 -3.04715  
PHE_3 -5.25196 0.29047 1.51945 0.02354 -0.38757 -1.99927  
VAL_4 -4.7212 0.38219 2.30519 0.01101 -0.33608 -4.69107  
ASP_5 -6.18239 0.68119 4.3013 0.02667 -1.21805 -4.77894  
LYS_6 -3.70129 0.51352 2.90682 0.00832 -0.90685 -3.25917  
ASN_7 -2.97985 0.42136 1.9979 0.00362 -0.17245 -2.08876  
GLY_8 -2.75353 0.28458 1.44949 1e-05 -0.20107 -4.00172  
ARG:CtermProteinFull_1294 -2.34298 0.24806 1.73197 0.01216 -0.06632 -0.34395
```

# Output Score Table

## Scores for individual terms:

```
# All scores below are weighted scores, not raw scores.
#BEGIN POSE ENERGIES TABLE dock 01 girk 1212 jon ML297 0500.pdb
label fa atr fa rep fa sol fa intra rep fa elec total
weights 0.8 0.4 0.6 0.004 0.42 NA
pose -5400.73 638.748 2428.12 13.5145 -302.036 -4267.34
GLN:NtermProteinFull_1 -1.02843 0.12113 0.83426 0.00595 -0.05882 0.54065
ARG_2 -5.46203 0.44148 4.02307 0.01303 -1.11204 0.00011
PHE_3 -5.25196 0.29047 1.51945 0.02354 -0.38757 -1.99927
VAL_4 -4.7212 0.38219 2.30519 0.01101 -0.33608 -4.69107
ASP_5 -6.18239 0.68119 4.3013 0.02667 -1.21805 -4.77894
LYS_6 -3.70129 0.51352 2.90682 0.00832 -0.90685 -3.25917
ASN_7 -2.97985 0.42136 1.9979 0.00362 -0.17245 -2.08876
GLY_8 -2.75353 0.28458 1.44949 1e-05 -0.20107 -4.00172
ARG:CtermProteinFull_1294 -2.34298 0.24806 1.73197 0.01216 -0.06632 -0.34395
```

**Total Score of Protein**

# Output Score Table

## Scores by amino acid position:

```
# All scores below are weighted scores, not raw scores.
#BEGIN POSE ENERGIES TABLE dock 01 girk 1212 jon ML297_0500.pdb
label fa atr fa rep fa sol fa intra rep fa elec total
weights 0.8 0.4 0.6 0.004 0.42 NA
pose -5400.73 638.748 2428.12 13.5145 -302.036 -4267.34
GLN:NtermProteinFull_1 -1.02843 0.12113 0.83426 0.00595 -0.05882 0.54065
ARG_2 -5.46203 0.44148 4.02307 0.01303 -1.13134 -3.04715
PHE_3 -5.25196 0.29047 1.51945 0.02354 -0.38757 -1.99927
VAL_4 -4.7212 0.38219 2.30519 0.01101 -0.33608 -4.69107
ASP_5 -6.18239 0.68119 4.3013 0.02667 -1.21805 -4.77894
LYS_6 -3.70129 0.51352 2.90682 0.00832 -0.90685 -3.25917
ASN_7 -2.97985 0.42136 1.9979 0.00362 -0.17245 -2.08876
GLY_8 -2.75353 0.28458 1.44949 1e-05 -0.20107 -4.00172
ARG:CtermProteinFull_1294 -2.34298 0.24806 1.73197 0.01216 -0.06632 -0.34395
```

More score term information

# Constraints (actually “restraints”)

Supplements energy function with additional information

- Commonly from experimental information

Various types:

atom\_pair\_constraint, dihedral\_constraint,  
angle\_constraint, coordinate\_constraint,  
residue\_type\_constraint ...


Interface with Rosetta using constraint files

# Additional score terms

Membrane terms:

fa\_mbsolv

fa\_mbenv



Over 100+ score terms

Yarov-Yarovoy, Schonbrun, and Baker 2006

Most are turned off (weight is set to 0), not in REF2015

To turn on score term, set weight to non-zero value

# Modifying Scorefunctions in Rosetta

## Modifying using the command line

1 - score:weights <filename>

2 - score:set\_weight <scoreterm<sub>1</sub>> <wt<sub>1</sub>> <scoreterm<sub>2</sub>> <wt<sub>2</sub>>

3 - score:patch <patchfile>

## Patchfile example

```
fa_atr = 0.423  
fa_rep = 0.100
```

# Modifying Scorefunctions in Rosetta

```
<ROSETTASCRIPTS>
  <SCOREFXNS>
    <ScoreFunction name="ligand_soft_rep" weights="ligand_soft_rep">
      <Reweight scoretype="fa_elec" weight="0.42"/>
    </ScoreFunction>
    <ScoreFunction name="hard_rep" weights="ligandprime">
    </ScoreFunction>
  </SCOREFXNS>
  <OUTPUT scorefxn="hard_rep" />
</ROSETTASCRIPTS>
```

## XML script options

- Weights filename or path to file
- Reweight specific terms as needed (as in patch file)
- Must include top-level output tag to ensure proper scoring in output files

# References:

## REF2015 References:

Alford RF, et. al, **The Rosetta All-Atom Energy Function for Macromolecular Modeling and Design.** *Journal of Chemical Theory and Computation*, **2017**. 13 (6), 3031-3048

Park H, et. al **Simultaneous Optimization of Biomolecular Energy Functions on Features from Small Molecules and Macromolecules** *Journal of Chemical Theory and Computation*, **2016**. 12 (12), 6201-6212

## Old Scorefxn (Talaris) References:

O'Meara MJ, et. al, **A Combined Covalent-Electrostatic Model of Hydrogen Bonding Improves Structure Prediction with Rosetta.** *Journal of Chemical Theory and Computation*, **2015**.

Leaver-Fay A, et. al **Scientific benchmarks for guiding macromolecular energy function improvement.** *Methods in enzymology*, **2013**. 523: p. 109.

# Useful Links

[https://www.rosettacommons.org/docs/latest/rosetta\\_basics/scoring/score-types](https://www.rosettacommons.org/docs/latest/rosetta_basics/scoring/score-types)

[https://www.rosettacommons.org/docs/latest/rosetta\\_basics/scoring/scoring-explained](https://www.rosettacommons.org/docs/latest/rosetta_basics/scoring/scoring-explained)